

A quasi-experimental study to assess the effect of a local anesthetic proparacaine on tropicamide-induced pupillary dilation

Jayashree Padmaraj Menishinkai, Chaitra Pujar, C. N. Manasa*, Laxman V. Dasar, Mallikarjun C. Salagar

Department of
Ophthalmology,
S. Nijalingappa Medical
College and Hanagal Shree
Kumareshwar Hospital,
Bagalkot, Karnataka, India

Received: 24 April 2015

Accepted: 11 May 2015

***Correspondence to:**

C. N. Manasa,
Email: manasacn17@gmail.
com

Copyright: © the author(s),
publisher and licensee Medip
Academy. This is an open-
access article distributed under
the terms of the Creative
Commons Attribution Non-
Commercial License, which
permits unrestricted non-
commercial use, distribution,
and reproduction in any
medium, provided the original
work is properly cited.

ABSTRACT

Background: Dilating the pupil is one of the most commonly practiced methods among ophthalmologists to examine the eye especially the posterior segment of the eye. Here we conducted a study to know the clinical effect of proparacaine, a local anesthetic, on tropicamide-induced pupillary dilation.

Methods: A Quasi-experimental study was conducted on 57 subjects aged between 20 and 70 years. Study was done over a period of 3 months (February 14 - April 14) at a tertiary health care center in Bagalkot after obtaining institutional ethical committee clearance. Proparacaine was instilled in one eye (right eye) before instilling tropicamide and the other eye with a placebo (normal saline). Pupil diameter was measured using a pupillary gauge at 0, 15, and 30 mins. Results were analyzed using unpaired Student's t-test.

Results: The mean clinically efficient pupillary diameter at the end of 15 mins in proparacaine instilled eye was 5.56 ± 0.5 and in control eye was 5.25 ± 0.45 mm with t value 8.13 ($p < 0.001$) which was statistically significant. And at the end of 30 mins pupillary diameter in the study group was 7.96 ± 0.43 mm and in the control group was 7.83 ± 0.43 mm with t value of 4.115 ($p < 0.001$) which was also statistically significant.

Conclusions: Therefore, we do recommend the use of proparacaine before instilling tropicamide for faster pupillary dilation.

Keywords: Proparacaine, Pupillary dilation, Tropicamide, Clinically effective diameter

INTRODUCTION

Dilation of pupils is an essential accompaniment to routine eye examinations, to facilitate the fundus evaluation using binocular indirect ophthalmoscope, fundus contact lenses, and other associated techniques.¹⁻³ It is essential not just for an ophthalmologist, but also for primary care practitioners.⁴ Pupillary dilation is very much essential in many intraocular surgeries like small incision cataract surgery, phacoemulsification, and outpatient procedures like laser treatments. A recommended drug regime for a mydriatic fundus examination uses the cholinergic antagonist tropicamide and the adrenergic agonist phenylephrine.⁵ The use of a topical anesthetic, namely proparacaine, before the instillation of any mydriatic has also been recommended earlier.⁵⁻¹¹ Mechanism of action is still not known, the corneal epithelium gets disrupted

when a topical anesthetic is used¹¹ and is thought to facilitate the absorption of the mydriatic agent through the cornea.^{5,12,13} Apart from that, prior instillation of anesthetic agent reduces reflex tearing, because of which a higher amount of mydriatic is available for corneal penetration.^{5,14} Advantages of using an anesthetic prior to instillation of a mydriatic consists of increased magnitude and rate of pupillary dilation,^{9,10,13} along with sustained pupillary dilation.¹⁰ Patient compliance will also be improved if a local anesthetic is used due to a reduction in the discomfort.^{5,14} In one of the study conducted by Ghose et al., it is reported that lignocaine has a significant increase in pupillary dilatation if it is used before instilling a drop of tropicamide indicating that prior instillation of a local anesthetic has a beneficial role to play in pupillary dilation.¹⁵

METHODS

This quasi-experimental study was conducted in the department of ophthalmology in the outpatient clinic from December 2014 to February 2015 over a period of 3 months. Informed written consent was taken from the individual participants who had taken part in the study. Institutional Ethics Committee clearance was obtained. Right eye was taken for the study while left eye served as control (Figure 1).

A total of 57 patients aged between 20 and 70 years were selected randomly by lot method attending outpatient clinic in our department. The subjects who were enrolled in the study underwent a thorough slit lamp examination with brief history taking. Subjects with pupillary abnormality, glaucoma, history of using any topical medication usage, trauma, pseudoexfoliation, uveitis, corneal scars, acute or chronic dacryocystitis, and any history of allergy to drugs were excluded from the study.

Undilated pupil measurements were taken from both eyes in a dark room using pupillary gauge.

Eye drops were instilled into the lower conjunctival fornix after asking the patient to look up toward the roof. And lacrimal passage was occluded by pressing on the medial side of the medial canthus so that drainage was reduced. Eye drops containing proparacaine was instilled in the study group and normal saline in the control group. After 5 mins tropicamide eye drops was instilled in both the groups.

Pupillary measurements were taken after 15 mins and at the end of 30 mins using the same pupillary gauge and the recordings were noted by the examiner. Instillation of the drop was done by one investigator and measurement of pupillary size at the end of 15 mins and 30 mins was done by another investigator separately.

Data were tabulated using Microsoft excel 2010 and analyzed using SPSS version 20 trial version. Student's



Figure 1: Pupillary gauge.

t-test was used to find the difference between two groups. For all practical purpose $p < 0.05$ was considered as statistically significant.

RESULTS

This study included 57 subjects, among which 22 (38.6%) were males and 35 (61.4%) were females. Among the 57 subjects, mean age was 58.42 ± 10.27 years. Right eye was the study eye and left eye was the control eye. Mean undilated pupil diameter (in mm) in a study group at 0 min was 2.89 ± 0.42 and in the control group was 2.89 ± 0.443 . There was not much difference in the pupillary diameter of both the study and the control group at 0 mins and it was not statistically significant. Six millimeter pupillary diameter (clinically effective diameter [CED]) is considered to be an adequate size for fundus evaluation, as reported from the previous study.¹⁶ The mean pupillary diameter in the study group at 15 mins was 5.56 ± 0.5 and in the control group was 5.25 ± 0.45 ($p < 0.001$) which was statistically significant. The pupillary diameter at the end of 30 mins in study group and control group was 7.96 ± 0.43 and 7.83 ± 0.43 , respectively ($p < 0.001$), which was statistically significant.

The pupillary size measurement done during the study was tabulated as follows.

The mean pupillary diameter in the right eye that is proparacaine treated eye increased from a mean pupillary diameter of 2.89 ± 0.42 at 0 min to 7.96 ± 0.43 at the end of 30 mins. On the other hand, mean pupillary diameter in the left eye that is placebo-treated eye increased from 2.89 ± 0.443 at 0 min to 7.83 ± 0.43 at the end of 30 mins. The difference in the mean pupillary diameter between the study eye and the control eye was statistically significant at 15 mins as well as 30 mins (Table 1).

Table 2 shows that 20 (35.1%) cases reached CED at the end of 15 mins and 37 (64.9%) cases did not reach CED in proparacaine treated eyes. In placebo-treated eye only 7 (12.3%) cases reached the CED and 50 (87.7%) cases did not reach CED at the end of 15 mins.

DISCUSSION

Instillation of one drop of proparacaine before the instillation of tropicamide for pupillary dilation resulted in a significant increase in pupillary diameter as compared to using tropicamide alone.

Ghose et al. studying the potentiating effect of 4% lignocaine on tropicamide-induced mydriasis obtained a mean maximum pupil size of 6.75 ± 0.80 mm in the intervention eye and 6.08 ± 0.97 mm in the control eye.¹⁵ This is much smaller than the pupil sizes of 7.96 ± 0.43 mm and 7.83 ± 0.43 mm, obtained for proparacaine-treated study

Table 1: Pupillary size in study eye (RE) and control eye (LE).

	Mean pupil dilatation (in mm)	SD	t value	p value
RE 0 min	2.895	0.4198	0.44	0.659
LE 0 min	2.886	0.4434		
RE 15 mins	5.561	0.5006	8.13	<0.001
LE 15 mins	5.246	0.4544		
RE 30 mins	7.965	0.4316	4.115	<0.001
LE 30 mins	7.833	0.4364		

SD: Standard deviation, RE: Right eye, LE: Left eye

Table 2: Proportion of study and control eyes reaching CED.

	CNED (%)	CED (%)
RE	37 (64.9)	20 (35.1)
LE	50 (87.7)	7 (12.3)
Total	87	27

CED: Clinically effective diameter, RE: Right eye, LE: Left eye

eyes and control eyes, respectively in the present study. The difference recorded in maximum pupil size, between both studies, was about 1.88 mm in the drug-treated eye and 1.75 mm in the control eye. The much larger difference in the drug-treated eye suggests that proparacaine may have a higher potentiating effect on tropicamide than eyes treated with lignocaine. It may be because of the lesser penetration and hence lesser availability of lignocaine as compared to proparacaine.

The mean pupillary diameter at the end of the study conducted by Ogun et al. was 6.01 ± 0.54 mm which is much less compared to the pupillary diameter in the proparacaine treated eye in our study which is 7.96 ± 0.43 at the end of our study. A typical Nigerian patient is dark skinned with densely pigmented iris. The difference in the pupillary diameter may be because of the increased pigmentation in a Nigerian subject as compared to ours.¹⁷ Same theory was suggested by Emiru, who states that iris pigmentation and not the race that is more likely to be responsible for the poorer pupil dilatation. Emiru noted that the dark-skinned African did not achieve the large pupil diameters even after 60 mins of dilatation.¹⁸ His reported mean pupillary diameter at 60 mins was 6.21 mm, which is much lesser to the mean diameter of 7.96 ± 0.43 mm obtained at 30 mins in proparacaine-treated eye in this study, which supports the theory of the higher pigmentation rate in African people as compared to Indian population here. Chen and Poth also could explain the reduced mydriatic effect due to decreased absorption in iris crypts because of obstruction by pigments within the chromophores.¹⁹

Topical anesthetics inhibit the rate of corneal epithelial cell migration by disrupting the cytoplasmic action in filaments and destroy superficial corneal epithelial microvilli.²⁰

The mechanism of action of potentiating effect of proparacaine on tropicamide-induced pupillary dilatation may be because of an increased intraocular penetration of tropicamide which is because of greater availability of the drug due to increased absorption through the microscopic disruptions in the corneal epithelium and reduced tearing because of which more time of contact and hence increased bioavailability might have been achieved.^{9,13,16,21,22}

CONCLUSION

It can be concluded that instilling proparacaine before tropicamide gives faster pupillary dilatation. Hence, we do recommend the use of proparacaine before instilling tropicamide for faster pupillary dilatation.

ACKNOWLEDGMENTS

Further, the authors wish to extend their gratitude to the anonymous reviewers for their valuable comments in improvising the research.

Funding: No funding sources

Conflict of interest: None declared

Ethical approval: The study was approved by the Institutional Ethics Committee

REFERENCES

- Phillips CI. Dilate the pupil and see the fundus. Br Med J (Clin Res Ed). 1984;288(6433):1779-80.
- Whitmer L. To see or not to see: routine pupillary dilatation. J Am Optom Assoc. 1989;60(7):496-500.
- Bartlett JR, Siderov J. Routine dilatation: appropriate choice of technique. Clin Exp Optom. 1995;78:38-9.
- Steinmann WC, Millstein ME, Sinclair SH. Pupillary dilatation with tropicamide 1% for funduscopic screening. A study of duration of action. Ann Intern Med. 1987;107(2):181-4.
- Talley DK, Bartlett JD. Topical and regional anesthesia. In: Bartlett JD, editor. Clinical Ocular Pharmacology. 3rd Edition. Boston: Butterworth-Heinemann; 1995: 463-78.
- Bartlett JD. Dilatation of the pupil. In: Bartlett JD, Jaanus SD, editors. Clinical Ocular Pharmacology. 3rd Edition. Boston: Butterworth-Heinemann; 1995: 479-502.
- Isenberg SW. The Eye in Infancy. 2nd Edition. St Louis: Mosby; 1994.
- Forman AR. A new low-concentration preparation for mydriasis and cycloplegia. Ophthalmology. 1980;87(3):213-5.
- Lyle WM, Bobier WR. Effects of topical anesthetics on phenylephrine-induced mydriasis. Am J Optom Physiol Opt. 1977;54(5):276-81.
- Mordi JA, Lyle WM, Mousa GY. Does prior instillation of

- a topical anesthetic enhance the effect of tropicamide? *Am J Optom Physiol Opt*. 1986;63(4):290-3.
11. Bryant JA. Local and topical anesthetics in ophthalmology. *Surv Ophthalmol*. 1969;13(5):263-83.
 12. Apt L, Henrick A. Pupillary dilatation with single eye drop mydriatic combinations. *Am J Ophthalmol*. 1980;89(4):553-9.
 13. Jauregui MJ, Polse KA. Mydriatic effect using phenylephrine and proparacaine. *Am J Optom Physiol Opt*. 1974;51(8):545-9.
 14. Kubo DJ, Wing TW, Polse KA, Jauregui MJ. Mydriatic effects using low concentrations of phenylephrine hydrochloride. *J Am Optom Assoc*. 1975;46(8):817-22.
 15. Ghose S, Garodia VK, Sachdev MS, Kumar H, Biswas NR, Pandey RM. Evaluation of potentiating effect of a drop of lignocaine on tropicamide-induced mydriasis. *Invest Ophthalmol Vis Sci*. 2001;42(7):1581-5.
 16. Pollack SL, Hunt JS, Polse KA. Dose-response effects of tropicamide HCl. *Am J Optom Physiol Opt*. 1981;58(5):361-6.
 17. Ogun OA, Oliver JW, Ashaye AO, Ajayi BG. Evaluating the potentiating effect of amethocaine on tropicamide-induced mydriasis in darkly pigmented irides, using infrared pupillometry. *Ophthalmol Eye Dis*. 2014;6:13-9.
 18. Emiru VP. Response to mydriatics in the African. *Br J Ophthalmol*. 1971;55(8):538-43.
 19. Chen KK, Poth EJ. Racial differences as illustrated by the mydriatic action of cocaine, euphthalmine and ephedrine. *J Pharmacol Exp Ther*. 1929;36:429-45.
 20. Brewitt H, Bonatz E, Honegger H. Morphological changes of the corneal epithelium after application of topical anaesthetic ointments. *Ophthalmologica*. 1980;180(4):198-206.
 21. Richardson RW. Comparing the mydriatic effect of tropicamide with respect to iris pigmentation. *J Am Optom Assoc*. 1982;53(11):885-7.
 22. Zauberman H, Neuman E. Tropicamide – a mydriatic cycloplegic in a West African eye clinic. *Cent Afr J Med*. 1970;16(1):8-11.

doi: 10.18203/2319-2003.ijbcp20150046

Cite this article as: Menishinkai JP, Pujar C, Manasa CN, Dasar LV, Salagar MC. A quasi-experimental study to assess the effect of a local anesthetic proparacaine on tropicamide-induced pupillary dilation. *Int J Basic Clin Pharmacol* 2015;4:410-3.