

## A prospective study of drug utilization pattern in cardiac intensive care unit at a tertiary care teaching hospital

H. Nagabushan<sup>1</sup>, H. S. Roopadevi<sup>1</sup>, G. M. Prakash<sup>2</sup>, R. Pankaja<sup>1\*</sup>

<sup>1</sup>Department of Pharmacology, Mandya Institute of Medical Sciences, Mandya, Karnataka, India, <sup>2</sup>Department of Medicine, Mandya Institute of Medical Sciences, Mandya, Karnataka, India

**Received:** 16 April 2015

**Accepted:** 11 May 2015

**\*Correspondence to:**

Dr. R. Pankaja,  
Email: pankaja.rudrappa@gmail.com

**Copyright:** © the author(s), publisher and licensee Medip Academy. This is an open-access article distributed under the terms of the Creative Commons Attribution Non-Commercial License, which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

### ABSTRACT

**Background:** Cardiovascular diseases remain the most common cause of sudden death. Appropriate drug therapy in cardiac intensive care unit (CICU) is crucial in managing cardiovascular emergencies and to decrease morbidity and mortality. The present study was conducted to observe the emergency cardiac diseases which are most frequently being treated and to study the prescribing prevalence among inpatients in CICU.

**Methods:** A prospective, observational study was carried out among 102 patients admitted in CICU at a tertiary care teaching hospital, Karnataka, for a period of 3 months. Demographic data, clinical history, and complete drug therapy received during their stay in CICU was noted.

**Results:** In our study, males (64.7%) had a higher incidence of cardiovascular emergencies than females (35.3%). Hypertension (32.4%) and Type 2 diabetes mellitus (28.4%) were the frequently associated co-morbid conditions. Antiplatelet drugs 80 (78.4%) was most commonly prescribed, followed by hypolipidemic drugs 75 (73.5%) and anticoagulants 65 (63.7%). The mean duration of stay in the hospital was 4.79±1.9 days. The average number of drugs per prescription was 7.8±2.2. Percentage of drugs prescribed by generic names was 52.9%. The percentage of drugs prescribed from essential drug list was 75.1%.

**Conclusions:** Antiplatelet drugs were the most frequently prescribed drug group. Mean number of drugs per prescription were high. The prescribing pattern could be improved by reducing the number of drugs per prescription and by prescribing generic drugs to reduce the economic burden of the patients.

**Keywords:** Cardiac intensive care unit, Cardiovascular diseases, Prescription

### INTRODUCTION

Cardiovascular diseases (CVDs) are the most frequent cause of morbidity and mortality throughout the world.<sup>1</sup> "Prescription" is defined as a written order, especially by a physician, for the preparation, and administration of a medicine or other treatment.<sup>2</sup>

Rational drug prescribing is defined as the use of the least number of drugs to obtain the best possible effect in the shortest period and at a reasonable cost.<sup>3</sup>

Polypharmacy is a major problem with cardiovascular inpatients admitted for a prolonged period of time.<sup>3</sup> Appropriate utilization of prescription drugs ensures favorable outcome of the patient and decreases mortality and morbidity.<sup>4</sup>

Studies on cardiac inpatients are rare and incomplete though risks of CVDs are frequently monitored. A prescription based survey is the most effective method to assess and evaluate the prescribing attitude of physicians.<sup>1</sup>

Hence, the present study aims at identifying the patient parameters, associated co-morbid conditions and the pattern of drug prescription among patients admitted in the cardiac intensive care unit (CICU) of a tertiary care teaching hospital.

The present study was conducted to observe the emergency cardiac diseases, which are most frequently being treated and to study the prescribing prevalence among inpatients in CICU.

## METHODS

This was a prospective, observational study carried out among patients admitted in CICU at a tertiary care teaching hospital, Karnataka, between months of November 2014 and January 2015.

The study was initiated after approval from the Institutional Ethics Committee. The data were collected from medical records of patients irrespective of their gender, age, and diagnosis. The collected data were analyzed with respect to demographic profile and drugs prescribed as following:

- Most common cardiovascular emergencies which are treated
- Drug related details: most common drug groups, dosage form, strength, route, frequency, duration of administration
- WHO criteria for prescription like average no of drugs per prescription, percentage of drugs prescribed by generic name, percentage of drugs with injections prescribed, and percentage of drugs prescribed from the WHO essential drugs list.

### Statistical analysis

Data were entered into Microsoft Excel (Windows 7; Version 2007) and analyses were done using the Statistical Package for Social Sciences (SPSS) for Windows software (version 18.0; SPSS Inc., Chicago). Descriptive statistics such as frequencies and percentages were calculated for categorical variables. Mean and standard deviation were computed for continuous variables. Graphic representation is used for visual interpretation of the analyzed data.

## RESULTS

The data of total 102 patients was obtained. Among them, males (64.7%) had a higher incidence of cardiovascular emergencies than females (35.3%). The mean age in years was 58.1±13.8 (Table 1). The incidence was higher in patients of age ≥55 years (65.7%).

Out of 102 patients, 61 (59.8%) had myocardial infarction (MI), 16 (15.7%) patients suffered from congestive heart failure (CHF), and the incidence of angina pectoris was 6 (5.9%) (Figure 1). Hypertension (32.4%) and Type 2 diabetes mellitus (28.4%) were the frequently associated co-morbid conditions.

Antiplatelet drugs 80 (78.4%) were most commonly prescribed, followed by hypolipidemic drugs 75 (73.5%), and anticoagulants 65 (63.7%) (Table 2).

Among antiplatelet drugs, aspirin 69 (67.6%), clopidogrel 59 (57.8%), and their combination 15 (14.7%) were prescribed, more often both drugs being preferred simultaneously. Streptokinase (100%) was the fibrinolytic

**Table 1: Socio-demographic profile of patients.**

Bio-social characteristics	Number	Percentage
Age (in years)		
≤35	12	11.8
36-45	9	8.8
46-55	14	13.7
56-65	38	37.3
>65	29	28.4
Mean±SD	58.1±13.8	
Range	20-80	
Gender		
Male	66	64.7
Female	36	35.3
Locality		
Urban	24	23.5
Rural	78	76.5
Marital status		
Married	96	94.1
Unmarried	6	5.9

SD: Standard deviation

**Table 2: Different categories of drugs prescribed to the patient.**

Treatment	Number	Percentage
Antiplatelets	80	78.4
Hypolipidemic	75	73.5
Anticoagulants	65	63.7
Drugs to reduce gastric acid secretion	57	55.9
ACE inhibitors	54	52.9
Antianginal drugs	50	49.0
Antibiotics	41	40.2
Beta blockers	39	38.2
Diuretics	36	35.3
Insulin and OHDs	34	33.3
CCBs	18	17.6
Inotropic drugs	14	13.7
Alpha blocker	08	07.8
Cardiac glycosides	08	07.8
Fibrinolytics	06	05.9
AT antagonists	05	04.9
Miscellaneous drugs	56	54.9


ACE: Angiotensin converting enzyme, OHD: Oral hypoglycemic drugs, CCBs: Calcium channel blockers

administered. Atorvastatin 73 (97.3%) and rosuvastatin (2.7%) were the hypolipidemic drugs prescribed. Low molecular weight heparin (100%) was the anticoagulant prescribed. Drugs prescribed to lower gastric acid secretion were pantoprazole 65 (84.4%) and ranitidine 12 (15.6%).

Among angiotensin converting enzyme (ACE) inhibitors, enalapril 33 (51.6%), and ramipril 28 (43.8%) were more commonly prescribed. Metoprolol 22 (56.4%), atenolol 12 (30.7%), and carvedilol 3 (7.7%) were the beta blockers prescribed. The calcium channel blockers (CCBs) prescribed were amlodipine 15 (83.3%) and diltiazem 3 (16.7%). Glyceryl trinitrate 47 (94.0%), Isosorbide dinitrate 4 (8.0%), and Isosorbide mononitrate 2 (4.0%) were the drugs prescribed under antianginals.

Furosemide 31 (75.0%) was the most common diuretic used, followed by spironolactone 3 (7.5%), mannitol 3 (7.5%), torasemide 2 (5.0%), and metolazone 2 (5.0%). Ceftriaxone 28 (68.3%), cefotaxime 10 (24.4%) were commonly prescribed drugs under antibiotic drug group. Dopamine 12 (85.7%) was the most common inotropic drug administered, followed by dobutamine 2 (14.3%). Digoxin (100%) was the prescribed cardiac glycoside for CHF. Among miscellaneous drug groups, analgesics (23.2%), methylxanthines (14.3%), and corticosteroids (12.5%) were routinely used.

The mean duration of stay in the hospital was 4.79±1.9 days. Table 3 gives the number of drugs prescribed per patient. The average number of drugs per prescription was 7.8±2.2. The



**Figure 1: Morbidity pattern of cardiovascular emergencies.**

**Table 3: Number of drugs prescribed per patient (n=102).**

Number of drugs prescribed per patient	Number	Percentage
4	04	03.9
5	03	02.9
6	16	15.7
7	20	19.6
8	31	30.4
9	17	16.7
10	08	07.8
11	01	01.0
20	02	02.0
Mean±SD	7.8±2.2	

SD: Standard deviation

dose, dosage form, route of administration, frequency, and duration were mentioned for 85.7%, 92.2%, 93.8%, 89.6% and 93.4% of drugs, respectively. Percentage of patients who were given injectable drugs was 100%. Percentage of drugs prescribed by generic names was 52.9% and percentage of brand names was 47.1%. The percentage of drugs prescribed from essential drug list was 75.1%.

**DISCUSSION**

The drug utilization study is been conducted widely and it is being carried out in different health care setups. Such studies are helpful to determine the behavior of the use of medicines in a society.<sup>5</sup> A survey based on prescription is considered to be one of the most cost effective methods to determine the prescribing approach of physicians.<sup>6</sup>

The present study is an attempt to evaluate the pattern of prescriptions of the patients admitted in CICU in a government set-up tertiary care teaching hospital.

A total of one hundred and two patient’s prescriptions were analyzed during 3 months period. Results indicated that male (64.7%) patients had a high frequency of cardiovascular incidences as compared to female (35.3%) patients which were in accordance with a study conducted by Al-Junid et al.<sup>7</sup> and Sreedevi et al.<sup>8</sup>

Out of 102 patients, 61 (59.8%) had MI, 16 (15.7%) patients suffered from CHF, and the incidence of angina pectoris was 6 (5.9%). Hypertension and diabetes were the two most common co-morbid conditions found in most of the patients, which were in accordance with study conducted by Dawalji et al.<sup>9</sup> and Christian et al.<sup>10</sup>

In a study conducted by Dawalji et al. it was found that most commonly prescribed drug classes were antiplatelet drugs followed by antihyperlipidemics.<sup>9</sup> Similar results were found in our study, as most commonly prescribed medication was Antiplatelet drugs 80 (78.4%), followed by hypolipidemic drugs 75 (73.5%). Among antiplatelet drugs, aspirin 69 (67.6%), clopidogrel 59 (57.8%), and their combination 15 (14.7%) were prescribed.

The beta-agonists such as dobutamine and dopamine are widely used to improve hemodynamics, in acutely decompensated chronic heart failure patients.<sup>11,12</sup> In our study, Dopamine 12 (85.7%) was the most common inotropic drug administered, followed by dobutamine 2 (14.3%).

A study was conducted by Muhit et al.<sup>13</sup> on cardiovascular disease prevalence and prescription patterns in Bangladesh. In this study among the lipid-lowering agents, atorvastatin was given to most of the patients whereas the second choice was rosuvastatin. Similarly in our study atorvastatin 73 (97.3%) was most commonly prescribed, followed by rosuvastatin (2.7%).

In the present study, low molecular weight heparin and streptokinase was the anticoagulant and fibrinolytic administered respectively.

The use of the ACE-inhibitors (52.9%) was higher than that of the  $\beta$ -blockers (38.2%) and the CCBs (17.6%). This was for keeping up with the increased use of ACE-inhibitors because of their beneficial effect on the morbidity and mortality in MI and ischemic heart disease patients, even after a long-term use. Furthermore, the ACE-inhibitors had a beneficial effect against fibrosis and cardiac remodelling.<sup>14</sup> These observations were in correlation with a study conducted by Pendhari et al., where ACE-inhibitors were most commonly prescribed compared to  $\beta$ -blockers and CCBs.<sup>1</sup>

Results showed that the average number of drugs prescribed in this study is  $7.8 \pm 2.2$ , which is comparable to studies conducted by Al-Junid et al. ( $7.56 \pm 3.37$ )<sup>7</sup> and Muhit et al. ( $6.35 \pm 1.56$ ).<sup>15</sup> Fixed-dose combinations (FDCs) are one of the strategies to reduce the number of drugs prescribed and to improve patient compliance. Increasing requirement of drugs when more than one disease is found in a patient justifies the use of FDCs.<sup>16,17</sup>

The overall percentage of generic drugs prescribed is 52.9%, which was found to be higher than the previous studies conducted by Al-Junid et al. (45.2%)<sup>7</sup>, Dawalji et al. (2.33%)<sup>9</sup>, and Tasneem et al. (6%).<sup>18</sup>

The average number of drugs in each prescription was  $7.8 \pm 2.2$ , percentage of drugs prescribed by generic name was 52.9%, percentage of injectables was 100%, and percentage of drugs prescribed from essential drug list was 75.1%, none of them were in compliance with WHO recommendations of  $\leq 2$ , 100%, 10%, and 70%,<sup>19,20</sup> respectively for each variable except for drugs prescribed by generic name.

Our study has several limitations; study duration was small, single centered with a small sample size. Since the study was done in a single government hospital, results of the study cannot be extrapolated to the general population.

The study would have been better if other regional hospitals of the city be included.

## CONCLUSION

Antiplatelet drugs followed by hypolipidemic drugs were the most frequently prescribed drug group. Mean number of drugs per prescription were high. The prescribing pattern could be improved by reducing the number of drugs per prescription and by prescribing generic drugs to reduce the economic burden of the patients as these drugs are used for long time. Prescription Indicators recommended by the WHO can help the Health Care Centers to obtain better organization and improve healthcare attention to the public.

## ACKNOWLEDGMENTS

We extend our gratitude to the staff, Department of General Medicine.

*Funding: No funding sources*

*Conflict of interest: None declared*

*Ethical approval: The study was approved by the Institutional Ethics Committee*

## REFERENCES

1. Pendhari SR, Chaudhari DR, Burute SR, Bite BM. A study on the drug utilization trends in the cardiovascular emergencies in a tertiary care hospital. *J Clin Diagn Res.* 2013;7(4):666-70.
2. Chandrashekar K, Saritha MK, Meti V. Prescription audit in tertiary care centre. *Asian J Med Clin Sci.* 2012;1(2):70-4.
3. Shankar R, Partha P, Shenoy N. Prescribing patterns of drugs among patients admitted with cardiovascular disorders in the internal medicine ward: prescribing patterns in inpatients. *Internet J Pharmacol.* 2001;1(2).
4. Barot PA, Malhotra SD, Rana DA, Patel VJ, Patel KP. Drug utilization in emergency medicine department at a tertiary care teaching hospital: a prospective study. *J Basic Clin Pharm.* 2013;4(4):78-81.
5. Introduction to Drug Utilization Research. World Health Organization 2003. Available at: <http://www.apps.who.int/medicinedocs/en/d/Js4876e/>. Accessed 08 April 2015.
6. Zafar F, Ali H, Naveed S, Korai OU, Rizvi M, Naqvi GR, et al. Drug utilization pattern in cardiovascular diseases: a descriptive study in tertiary care settings in Pakistan. *J Bioequiv Availab.* 2015;7(1):059-62.
7. Al-Junid SM, Ezat WP, Surianti S. Prescribing patterns and drug cost among cardiovascular patients in Hospital Universiti Kebangsaan Malaysia. *Med J Malaysia.* 2007;62(1):59-65.
8. Sreedevi K, Rao JV, Fareedullah M, Vijayakumar S. A study on prescription pattern of statins in cardiovascular disease. *Der Pharm Lett.* 2011;3(3):393-6.
9. Dawalji S, Venkateshwarlu K, Thota S, Venisetty PK, Venisetty RK. Prescribing pattern in coronary artery disease: a prospective study. *Int J Pharm Res Rev.* 2014;3(3):24-33.
10. Christian RP, Rana DA, Malhotra SD, Patel VJ. Evaluation of rationality in prescribing, adherence to treatment guidelines, and direct cost of treatment in intensive cardiac care unit: a prospective observational study. *Indian J Crit Care Med.* 2014;18(5):278-84.
11. Parissis JT, Farmakis D, Nieminen M. Classical inotropes and new cardiac enhancers. *Heart Fail Rev.* 2007;12(2):149-56.
12. Sivasankar V, Ramalakshmi S, Harika T, Ramesh M, Kousalya K. The prescribing pattern of dopamine and dobutamine in congestive heart failure patients in a tertiary care teaching hospital. *World J Pharm Sci.* 2014;2(4):404-8.
13. Muhit MA, Rahman MO, Raihan SZ, Asaduzzaman M, Akbar MA, Sharmin N. Cardiovascular disease prevalence and prescription patterns at a tertiary level hospital in Bangladesh. *J Appl Pharm Sci.* 2012;2(3):80-4.
14. Tripathi KD. Drug affecting renin-angiotensin system and plasma Kinins. In: Tripathi KD, editor. *Text Book of Essentials of Medical Pharmacology.* 7th Edition. New Delhi: Jaypee Brother Medical Publication; 2013: 495-511.

15. Patel MK, Barvaliya MJ, Patel TK, Tripathi C. Drug utilization pattern in critical care unit in a tertiary care teaching hospital in India. *Int J Crit Illn Inj Sci*. 2013;3(4):250-5.
16. Bangalore S, Kamalakkannan G, Parkar S, Messerli FH. Fixed-dose combinations improve medication compliance: a meta-analysis. *Am J Med*. 2007;120(8):713-9.
17. Rathnakar UP, Shenoy A, Ullal SD, Shivaprakash, Sudhakar P, Shastry R. Prescribing patterns of fixed dose combinations in hypertension, diabetes mellitus and dyslipidemia among patients attending a cardiology clinic in a tertiary care teaching hospital in India. *Pharm Glob (IJCP)*. 2011;2(6):1-3.
18. Tasneem S, Fouzia N. Drug utilization study in ischemic heart diseases associated with diabetes and hypertension. *Int J Pharm Bio Sci*. 2010;1(3):1-4.
19. Vooss AT, Diefenthaler HS. Evaluation of prescription indicators established by the WHO in Getulio Vargas - RS. *Braz J Pharm Sci*. 2011;47(2):385-90.
20. Vries TP, Henning RH, Hogerzeil H, Fresle DA. WHO Guide to Good Prescribing. Available at: <http://www.apps.who.int/medicinedocs/pdf/whozip23e/whozip23e.pdf>. Accessed 08 April 2015.

**doi:** 10.18203/2319-2003.ijbcp20150044

**Cite this article as:** Nagabushan H, Roopadevi HS, Prakash GM, Pankaja R. A prospective study of drug utilization pattern in cardiac intensive care unit at a tertiary care teaching hospital. *Int J Basic Clin Pharmacol* 2015;4:579-83.